AD	

Award Number: DAMD17-99-1-9142

TITLE: Mechanism of Growth Factor Attenuation of Cell Death in

Chemotheraphy Treated Breast Cancer Cells

PRINCIPAL INVESTIGATOR: Carla VanDenBerg

CONTRACTING ORGANIZATION: University of Colorado Health Sciences

Center

Denver, Colorado 80262

REPORT DATE: August 2001

TYPE OF REPORT: Annual Summary

PREPARED FOR: U.S. Army Medical Research and Materiel Command

Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT:

Approved for public release; distribution unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

20020416 133

REPORT DOCUMENTATION PAGE

Form Approved OMB No. 074-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of

Management and Budget, Paperwork Reduction Proje	ct (0704-0188), Washington, DC 20503		
1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE	•	E AND DATES COVERED
	August 2001	Annual Sum	mary (1 Aug 00 - 31 Jul 01)
4. TITLE AND SUBTITLE			5. FUNDING NUMBERS
Mechanism of Growth Fact	DAMD17-99-1-9142		
Chemotheraphy Treated Breast Cancer Cells.			1
6. AUTHOR(S)			
Carla VanDenBerg			
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)			8. PERFORMING ORGANIZATION
	3-1-1- 0-1		REPORT NUMBER
University of Colorado Health Sciences			
Center			
Denver, Colorado 80262			
E-Mail:			
9. SPONSORING / MONITORING AGE	NCY NAME(S) AND ADDRESS(ES)	10. SPONSORING / MONITORING
			AGENCY REPORT NUMBER
U.S. Army Medical Research and M			
Fort Detrick, Maryland 21702-5012			
11. SUPPLEMENTARY NOTES			
12a. DISTRIBUTION / AVAILABILITY STATEMENT			12b. DISTRIBUTION CODE
,			
Approved for public release; distribution unlimited			
		v v	
13. ABSTRACT (Maximum 200 Words	;/		-
1			

Upregulation of IGF-I (Insulin-like Growth Factor) receptor dependent pathways like phosphatidylinositol 3-kinase (PI 3-kinase) may diminish chemosensitivity of breast cancer cells via c-Jun N-terminal Kinase (JNK) response. In this project we have determined that IGF-I activates the survival pathway involving PI 3-kinase and Akt. Using stable transfection approaches we have confirmed the survival effect of activated Akt in MCF-7 cells. Surprisingly, we have also observed that IGF-I induces JNK in breast cancer cells. IGF-I and chemotherapy co-treatment results in additive effect on JNK activity, in contrast to our predicted results that IGF-I would antagonize chemotherapy induced JNK activity. PI 3-kinase mediates IGF-I induction of JNK in MCF-7 cells but not in T47D cells. We plan to further confirm the importance of JNK in IGF-I mediated survival effects in breast cancer cells.

14. SUBJECT TERMS	15. NUMBER OF PAGES		
IGF-I, chemotherapy, s	5		
			16. PRICE CODE
17. SECURITY CLASSIFICATION	18. SECURITY CLASSIFICATION	19. SECURITY CLASSIFICATION	20. LIMITATION OF ABSTRACT
OF REPORT	OF THIS PAGE	OF ABSTRACT	
Unclassified	Unclassified	Unclassified	Unlimited

Table of Contents

Cover 1	
SF 298	<u>.</u>
Table of Contents ³	}
Introduction	ŀ
Body	<u>'</u>
Key Research Accomplishments5	5
Reportable Outcomes5	5
Conclusions5	5
References	/ A
Appendices _N .	/Δ

Introduction

Cancer patients experience several toxicities from chemotherapy. One frequent toxicity is peripheral neuropathy, and strategies to relieve chemotherapy-induced neuropathy have had little success. Clinical trials using IGF-I have been proposed to combat chemotherapy-related neurotoxicities. IGF-I has trophic effects on motor neurons and is used for treatment of amyotrophic lateral sclerosis (ALS or Lou Gerig's disease) and diabetic neuropathy. However, there is concern about the potential for IGF-I to stimulate tumor growth or inhibit chemotherapy effects in patients receiving various treatments.

Body

Research accomplishments: We are studying the effects of IGF-I on breast cancer cells, focusing particularly on its ability to inhibit chemotherapy activity. We hypothesized that IGF-I acts through its receptor to induce at least one cell survival pathway. Task 1 (to characterize IGF-I receptor cytoprotective effects on PI 3-kinase) of this project was essentially completed in Year 1 of funding, as discussed in previous progress report.

Task 2 of this project is to characterize the potential interactions between the PI 3-kinase survival and JNK stress-induced signaling pathways by exposure to IGF-I -/+ chemotherapy using pharmacologic and molecular techniques to confirm these interactions. Previously, we had shown that JNK activity is induced by Taxol and Taxotere treatment in MCf-7 cells. Surprisingly, IGF-I treatment of cells markedly enhanced JNK activity as well. Co-culture with a PI 3-kinase inhibitor reduced IGF-I mediated JNK activity. In addition, when cells were cotreated with IGF-I and chemotherapy, JNK response was enhanced by co-treatment, rather than abrogated by IGF-I, as we had initially predicted. In order to better delineate the function of JNK in our model, MCF-7 cells were transfected with wildtype and dominant negative forms of JNK. Stable transfectants were cultured in Taxol alone, IGF-I alone, or Taxol + IGF-I. Our results generally support a role for JNK in mediating cell death. Our data do not support a role for JNK3 in mediating IGF-I survival responses, but we may not have observed this interaction for various reasons.

In the past year, we have also been studying if IGF-I induction of JNK may be observed in other cell lines especially since other investigators have published that JNK may confer survival effects only in cell expressing mutant p53. Since MCF-7 cells express wild-type p53, we decided to study the T47D breast cancer cell line. Indeed, we observe IGF-I induction of JNK in these breast cancer cells as well. In contrast to MCF-7 cells, IGF-I induction of JNK is not dependent upon PI 3-kinase. We have also developed a mutant p53 overexpressing transfectant of the MCF-7 cells to study JNK survival effects more completely in the future. We plan to further characterize the importance of p53 and PI 3-kinase in JNK function.

In reconsideration of the other proposed work in Task 2, to identify substrate specificity of JNK induced by chemotherapy or IGF-I, we believe we can more directly address the role of IGF-I activated JNK using JNK antisense approaches. We are establishing a collaboration to develop antisense approaches to JNK isoforms to ultimately determine if IGF-I survival effects are mediated in part through JNK.

We have initiated studies included in Task 3 which is to characterize if IGF-I and chemotherapy interactions in breast cancer cell lines are p53-dependent and compare ER positive and ER negative breast cancer cells with an immortalized, noncancerous breast epithelial cell line. These studies are partially completed, as discussed above in Task 2. We will continue to characterize additional breast cell lines.

Training accomplishments: In the past several months we have been developing an ecdysone inducible expression system in our MCF-7 cells in order to more clearly identify the roles of Akt and JNK in our model. Dr. Horwitz's laboratory has successfully developed an ecdysone-inducible system in the T47D cells and has been advising us in optimizing our conditions.

Reviewer comments on our manuscript submitted to *Cancer Research* requested two important areas for additional data to accept our manuscript. First, we needed to demonstrate if our observations in MCF-7 cells could be applied to other cell lines. We believe we have done this using the T47D cells. Secondly, the reviewer wanted an additional apoptosis assay to confirm our initial work. This latter request has required extensive effort to identify an assay that may be used to measure apoptosis in MCF-7 cells. MCF-7 cells lack expression of caspase 3, a protein responsible for many of the morphological characteristics of cell apoptosis. We believe we have finally accomplished this goal and plan to resubmit our manuscript shortly.

Key Research Accomplishments:

- Additional apoptosis assay development
- In contrast to MCF-7 cells, IGF-I induction of JNK is PI 3-kinase independent in T47D cells
- Similar to MCF-7 cells, IGF-I and stress treatments of breast cancer cells lead to increased JNK activity compared to either treatment alone

Reportable Outcomes

Manuscripts

Mamay, C.L., Wolf, D.M., Molina, D.M., Van Den Berg, C.L. "The cooperative and opposing roles of JNK and Akt in IGF-I mediated survival of chemotherapy treated breast cancer cells". Submitted to *Cancer Research*.

Funding Based on Work Supported by this Award

Grants Funded:

"IGF-I survival effects on p53 induced apoptosis." Charlotte Geyer Foundation, November 2000 to May 2001. Total budget \$58,000 (bridge funding mechanism).

"IGF-I survival effects on p53 induced apoptosis." National Cancer Institute, June 2001 to July 2004. Principal Investigator. Total Budget: \$509,625.